

University of Groningen

Over de behandeling van patienten met hypospadie

Heybroek, Gerard

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:

1964

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Heybroek, G. (1964). *Over de behandeling van patienten met hypospadie*. [, Rijksuniversiteit Groningen]. [S.n.].

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

STELLINGEN

- i. Bij de behandeling van patiënten met hypospadie dient men ernaar te streven deze behandeling tegen de schoolleeftijd te hebben voltooid.
- ii. De reconstructie der perifere urethra bij een patiënt, lijdende aan hypospadie, dient pas te geschieden als men de zekerheid heeft dat er geen ventrale curvatuur aanwezig is.
- iii. Indien bij een geval van hypospadie het orificium urethrae externum zich in de buurt van de peno-scrotale hoek bevindt, verdient de reconstructiemethode volgens CECIL de voorkeur wegens de geringe kans op fistelvorming.
- iv. Hypospadie is te beschouwen als een vorm van manlijk pseudo-hermaphroditisme.
- v. Het vroeg sluiten van een myelomeningocele heeft geen invloed op de neurologische uitvalsverschijnselen.
- vi. Wanneer een algemeen ziekenhuis beschikt over een zogenaamde revalidatieafdeling, mag als onderdeel daarvan een arbeidstherapie-afdeling niet ontbreken.
- vii. Voor het vroegtijdig vaststellen van chloroquinebeschadiging van de retina is een E.O.G. onontbeerlijk.

- viii. Het verdient aanbeveling bij de chirurgische behandeling van de cheilognathopalatoschisis prophylactisch antibiotica toe te dienen.

R. M. A. Mc. Clelland and T. J. S. Patterson Brit. J. Plast. Surg. 16 : 144 (1963).

- ix. De toediening van penicilline en chloramfenicol in hoge doses vermindert bij kinderen het optreden van neurologische restverschijnselen na acute etterige meningitis.
- x. Een behandelingswijze, waarvan de waarde nog niet statistisch is bewezen, dient niet om die reden te worden afgewezen.

Stellingen behorende bij G. Heybroek
Over de behandeling van patiënten met hypospadie
Groningen 1964